

Neck Lift

Performed either alone or as part of a face lift, a neck lift can remove excess skin, fatty tissue and banding, and restore a smooth, refined appearance to the neck.

About Neck Lift Surgery

For people with loose sagging skin under the chin, vertical bands in the neck, "turkey wattles" or necks that are too heavy, neck lift surgery – either alone or in combination with other cosmetic procedures – can produce very gratifying results.

Different procedures are used to address each of these problems:

- Sagging skin may be tightened or removed using a procedure called Cervicoplasty.
- To correct vertical bands or wattles, a technique called Platysmaplasty can be used to tighten the muscles of the neck.
- Fat deposits can be removed directly or by liposuction.

The quality of the result will depend upon the elasticity of the neck skin and degree of sun damage to the area. Naturally, it is important to have realistic expectations, but fortunately current techniques allow for an elaborate and detailed restitution of a youthful contouring the neck, that can benefit patients with a variety of problems.

In general terms, people whose relatives have strong jaws and chins, maintain a more defined appearance of the neck later in life. For those who have inherited a relatively small chin, a combination of neck lift and chin augmentation can produce dramatic improvements for the facial balance and neck contours.


A relatively small group of people tend to have poor definition in the neck very early in life, without ever being overweight. When the elasticity of the skin of the neck is good, it is possible to do an extensive contouring, through an approach called anterior neck lift, which uses only a 3 cm incision under the chin.

- Through this access it is possible to reduce the fatty tissue situated below the skin as well as the one deep to the platysma muscles, trim the lower pole of the submandibular glands if required, and adjusts the position and tension of the platysma muscles, in order to optimise the neck contour.
- It is by necessity a more powerful approach and then simple liposuction, which can only address the fatty tissue situated under the skin.

Your consultation

During your consultation your surgeon will take a general medical history and ask about other surgeries you may have had. Then the two of you will discuss your goals for the procedure and your surgeon will explain what results you can reasonably expect. Often, in particular for patients after their mid-40s, the lower part of the submandibular glands is palpable below the jaw line, and with a more simplistic approach to neck rejuvenation may sometimes become more visible at the surgery. It is possible during the neck lift procedure to address both the superficial and deep seated fatty tissue in the neck, as well as trim the lower poles of the submandibular glands, for a more detailed neck contouring.

You'll decide which procedure or combination of procedures is best and whether you should have your neck lift in conjunction with a face lift or other cosmetic procedure. You will also decide what kind of anaesthesia will be used.



15 Railway Road
Subiaco 6008
Western Australia

info@drbriggs.com.au
T 08 9381 9300
F 08 9381 9299

Patrick Briggs
Plastic Surgeon Pty Ltd
ABN 80 084 350 274

Member of the
Australian Society
of Plastic Surgeons

Member of the
Australasian Society of
Aesthetic Plastic
Surgeons

Member of the
International Society of
Plastic Surgeons



Our services

From your first visit to Dr Briggs' Subiaco office, you'll work directly with Consultant Plastic, Cosmetic and Reconstructive Surgeon Mr. Patrick Briggs, the practice's director.

Together you will explore the possibilities and discuss the results you can expect.

Imaging software will be used to show you how you might look after your procedure and every aspect of your treatment, from evaluation, through surgery to post surgical care will be carefully explained.

A very personal choice

A subtle blend of art and science, cosmetic surgery has become increasingly safe and comfortable in recent years.

Techniques have been refined, new materials have been developed and results have become more natural, predictable and satisfying. No other kind of surgery allows the surgeon more creativity; and no other kind of surgery is more deeply personal for the patient.





Your surgery & what to expect

Neck lift surgery usually takes between two and three hours, longer if it is being performed in conjunction with a face lift or other procedure. If it is being performed alone, neck lift surgery is usually done as a day case.

- If you have vertical bands or wattles that need to be addressed by platysmaplasty surgery, your surgeon will make small incisions under your chin or behind your ears to reach the your neck muscles and move them as needed.
- If you are having cervicoplasty surgery to correct loose and sagging skin, your surgeon will make similar incisions, trim the skin, move it into place and secure it using stitches or tissue glue. Your surgeon may also fit you with a compression bandage and instruct you to wear it continuously for at least a week.
- If excess fat must be removed, you surgeon may use liposuction, a process whereby fat is vacuumed away through thin cannulas. If this is the only procedure you are having it should take an hour or less. For a more extensive contouring, even in the presence of excess fatty tissue which can be related to genetic predisposition, lipodystrophy or excess weight, direct reduction of the fatty tissue through an incision under the chin is the procedure of choice.

After your procedure

In the days and weeks after your surgery there are several things you should expect:

- Depending on how extensive your procedure was, you may have some swelling and bruising. Both should settle in four to ten days.
- You may also experience feelings of tightness, burning, pulling and numbness. These are normal in the first few weeks and will resolve on their own.
- If you have sutures that need removing they will be taken out a week to ten days after your surgery.
- Any discomfort you experience can be controlled by taking paracetamol or medication provided by the hospital. Do not take aspirin or other anti-inflammatory medication as these can promote bleeding.
- If you smoke, you will need to stop entirely for two weeks before and two to three weeks after your surgery to allow your body to heal in the best conditions.
- You should be able to return to work in one to two weeks, however you will need to avoid physical activities like sports for at least three weeks, possibly longer.
- Firmness of the neck tissues is common after having a Neck Lift. It may take some weeks to settle.

Mr. Patrick Briggs

Mr. Patrick Briggs is an internationally trained plastic and cosmetic surgeon based in Subiaco, Western Australia.

After graduating from the Royal College of Surgeons in 1977, Mr. Briggs trained in general surgery at St. Bartholomew's Hospital in London, obtaining the fellowship Examination of the Royal College of Surgeons of England in May 1981 and the Fellowship Examination of the Royal College of Surgeons in Ireland in June 1981.

Training in Plastic Surgery followed at Mount Vernon Hospital and University College Hospitals in London, Allegheny General Hospital in Pittsburgh, Pennsylvania, USA and Cork University Hospital, Ireland.

In 1991, Mr. Briggs passed the Intercollegiate Board Examination in plastic surgery as administered by the four Royal Colleges of Great Britain and Ireland – the FRCS (Plastic Surgery). That same year, he was awarded a Certificate of Completion of Higher Surgical Training (CCST) in plastic and cosmetic surgery.

15 Railway Road
Subiaco 6008
Western Australia

info@drbriggs.com.au
T 08 9381 9300
F 08 9381 9299

Patrick Briggs
Plastic Surgeon Pty Ltd
ABN 80 084 350 274

Member of the
Australian Society
of Plastic Surgeons

Member of the
Australasian Society of
Aesthetic Plastic
Surgeons

Member of the
International Society of
Plastic Surgeons

